

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior School Certificate Examination, 2024-25
SUBJECT NAME CHEMISTRY (Theory) -043
(Q.P.CODE 56/7/2) MM: 70

General Instructions: -

You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.

“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its’ leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under various rules of the Board and IPC.”

Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one's own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. **However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-X, while evaluating two competency-based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.**

The Marking scheme carries only suggested value points for the answers

These are in the nature of Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.

The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.

Evaluators will mark(✓) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives an impression that answer is correct and no marks are awarded. **This is most common mistake which evaluators are committing.**

If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.

If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.

If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out with a note **“Extra Question”**.

No marks to be deducted for the cumulative effect of an error. It should be penalized only once.

A full scale of marks _____(example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.

Every examiner has to necessarily do evaluation work for full working hours i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.

Ensure that you do not make the following common types of errors committed by the Examiner in the past:-

- Leaving answer or part thereof unassessed in an answer book.
- Giving more marks for an answer than assigned to it.
- Wrong totaling of marks awarded on an answer.
- Wrong transfer of marks from the inside pages of the answer book to the title page.
- Wrong question wise totaling on the title page.
- Wrong totaling of marks of the two columns on the title page.
- Wrong grand total.
- Marks in words and figures not tallying/not same.
- Wrong transfer of marks from the answer book to online award list.
- Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.)

Half or a part of answer marked correct and the rest as wrong, but no marks awarded.


While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.

Any unassessed portion, non-carrying over of marks to the title page, or totaling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.

The Examiners should acquaint themselves with the guidelines given in the “**Guidelines for Spot Evaluation**” before starting the actual evaluation.

Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.

The candidates are entitled to obtain photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

Q.No	Value points	Mark										
SECTION A												
1	C	1										
2	D	1										
3	B	1										
4	C	1										
5	B	1										
6	A	1										
7	B	1										
8	A	1										
9	C	1										
10	D	1										
11	A	1										
12	C	1										
13	A	1										
14	C	1										
15	A	1										
16	A	1										
SECTION B												
17	When vapour pressure of the solution is lower than expected from the ideal behaviour. Example : chloroform and acetone/ HNO_3 and water (or any other suitable example) Maximum boiling azeotrope	1 $\frac{1}{2}$ $\frac{1}{2}$										
18	a) Dichloridobis(ethane-1,2-diamine)platinum(IV) ion 	1 1										
OR												
18	i) $[\text{Co}(\text{NH}_3)_5(\text{CO}_3)]\text{Cl}$ ii) Pentaamminenitrito-O-cobalt(III) chloride	1 1										
19	When one of the reactant is present in excess Hydrolysis of an ester/ sucrose (or any other suitable example) For elementary reaction, which takes place in a single step.	1 $\frac{1}{2}$ $\frac{1}{2}$										
20	<table><tr><th>Amylose</th><th>Amylopectin</th></tr><tr><td>Linear polymer of α-glucose</td><td>Branched polymer of α-glucose</td></tr><tr><td>C_1-C_4 glycosidic linkage</td><td>C_1-C_4 and C_1-C_6 glycosidic linkages</td></tr><tr><td>Water soluble</td><td>Water insoluble</td></tr><tr><td>It constitutes 15-20% of starch</td><td>It constitutes 80-85% of starch</td></tr></table> <p style="text-align: right;">(Any TWO)</p>	Amylose	Amylopectin	Linear polymer of α -glucose	Branched polymer of α -glucose	C_1 - C_4 glycosidic linkage	C_1 - C_4 and C_1 - C_6 glycosidic linkages	Water soluble	Water insoluble	It constitutes 15-20% of starch	It constitutes 80-85% of starch	1+1
Amylose	Amylopectin											
Linear polymer of α -glucose	Branched polymer of α -glucose											
C_1 - C_4 glycosidic linkage	C_1 - C_4 and C_1 - C_6 glycosidic linkages											
Water soluble	Water insoluble											
It constitutes 15-20% of starch	It constitutes 80-85% of starch											
21	-Because C—X bond acquires a partial double bond character due to resonance/ sp^2 hybridized carbon of C-X bond leading to shorter bond length (Or any other suitable reason). -Nitro group withdraws the electron density from the benzene ring and thus facilitates the attack of the nucleophile on haloarene / $-\text{NO}_2$ group being electron withdrawing stabilises the intermediate carbanion.	1 1										

	SECTION C	
22	<p>a) The amino acids, which can be synthesised in the body, are known as non- essential amino acids whereas those which cannot be synthesised in the body and must be obtained through diet, are known as essential amino acids.</p> <p>b) A linkage which joins two amino acids through -CO-NH- bond is peptide linkage whereas a linkage which joins two monosaccharides through oxygen atom is known as glycosidic linkage.</p> <p>c) Fibrous protein: parallel polypeptide chain structure / insoluble in water Globular protein: spherical polypeptide chain structure/ soluble in water</p>	<p>1</p> <p>1</p> <p>1</p>
23	<p>a) Because the carboxyl group is deactivating and the catalyst aluminium chloride (Lewis acid) gets bonded to the carboxyl group.(forms salt)</p> <p>b) Because carbonyl carbon of HCHO is more electrophilic than CH₃CHO/ due to +I effect of methyl group/ steric effect of methyl group, CH₃CHO is less reactive.</p> <p>c) Because of greater electronegativity of sp² hybridised carbon to which carboxyl carbon is attached.</p>	<p>1</p> <p>1</p> <p>1</p>
24	<p>a) <chem>CH3I</chem> + <chem>c1ccccc1O</chem></p> <p>b) <chem>Oc1c([N+](=O)[O-])cc([N+](=O)[O-])cc1[N+](=O)[O-]</chem></p> <p>c) <chem>OC1CCCCC1</chem></p>	<p>1</p> <p>1</p> <p>1</p>
25	$ \begin{array}{ccccc} & 2\text{N}_2\text{O}_5(\text{g}) & \rightarrow & 2\text{N}_2\text{O}_4(\text{g}) & + & \text{O}_2(\text{g}) \\ \text{Start } t = 0 & p_i \text{ atm} & & 0 \text{ atm} & & 0 \text{ atm} \\ \text{At time } t & (p_i - 2x) \text{ atm} & & 2x \text{ atm} & & x \text{ atm} \end{array} $ <p> $P_t = P_i - 2x + 2x + x = P_i + x$ $x = P_t - P_i$ $p_A = P_i - 2x$ $= P_i - 2(P_t - P_i)$ $= 3P_i - 2P_t$ </p> <p> $k = \frac{2.303}{t} \log \frac{P_i}{p_A}$ </p> <p>Where $p_i = 0.5 \text{ atm}$, $p_A = 3p_i - 2p_t$ $= (3 \times 0.5) - (2 \times 0.625)$ $= 0.25 \text{ atm}$ </p> <p> $k = \frac{2.303}{100 \text{ s}} \log \frac{0.5 \text{ atm}}{0.25 \text{ atm}}$ </p> <p> $= \frac{2.303}{100 \text{ s}} \times 0.3010$ $= 6.93 \times 10^{-3} \text{ s}^{-1}$ </p>	<p>1</p> <p>1</p> <p>1</p>

26.	<p>A=</p> $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - [\\ \\ \text{CH}_3 \end{array}$ <p>B=</p> $\begin{array}{c} \text{CH}_3 - \text{C} = \text{CH}_2 \\ \\ \text{CH}_3 \end{array}$ <p>C=</p> $\begin{array}{c} \text{I} \\ \\ \text{CH}_3 - \text{CH} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$ <p>D=</p> $\begin{array}{c} \text{CH}_3\text{CHCH}_2\text{CH}_2\text{CHCH}_3 \\ \qquad \qquad \\ \text{CH}_3 \qquad \qquad \text{CH}_3 \end{array}$ <div style="border: 1px solid black; padding: 10px; margin-top: 10px;"> $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{I} \\ \\ \text{CH}_3 \end{array} \xrightarrow[\text{KOH(aq)/}\Delta]{\text{KOH(aq)/}\Delta} \begin{array}{c} \text{CH}_3 - \text{C} = \text{CH}_2 \\ \\ \text{CH}_3 \end{array} + \text{KI} + \text{H}_2\text{O}$ </div>	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p>
27	<p>(a) Lead storage battery</p> <p>Anode: $\text{Pb(s)} + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{PbSO}_4(\text{s}) + 2\text{e}^-$</p> <p>Cathode: $\text{PbO}_2(\text{s}) + \text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O(l)}$</p>	<p>1</p> <p>1</p> <p>1</p>
OR		
27	<p>(b) Because at cathode the reaction with higher value of E° is preferred and therefore, the reduction of H_2O to H_2 gas is preferred whereas at anode water should get oxidised in preference to $\text{Cl}^-(\text{aq})$, however, on account of overpotential of oxygen, oxidation of Cl^- to Cl_2 gas is preferred.</p> <p>$\text{NaCl(aq)} + \text{H}_2\text{O(l)} \rightarrow \text{Na}^+(\text{aq}) + \text{OH}^-(\text{aq}) + \frac{1}{2}\text{H}_2(\text{g}) + \frac{1}{2}\text{Cl}_2(\text{g})$</p>	<p>1</p> <p>1</p> <p>1</p>
28	<p>$\Delta T_f = iK_f m$</p> $\Delta T_f = \frac{i \times K_f \times w_2 \times 1000}{M_2 \times w_1}$ $0.45 = \frac{i \times 5.12 \times 0.3 \times 1000}{60 \times 30}$ <p>$i = 0.527$</p> $\alpha = \frac{i-1}{1/n-1}$ $\alpha = \frac{0.527-1}{1/2-1} \quad (n=2)$ <p>$\alpha = 0.946$ or 94.6% (Or any other suitable method)</p>	<p>½</p> <p>1</p> <p>½</p> <p>½</p> <p>½</p>

	SECTION D	
29	<p>a) (i)</p> <div style="border: 1px solid black; padding: 10px; margin: 10px auto; width: fit-content;"> $\text{CH}_3\text{COOH} \xrightarrow{\text{NH}_3, \text{Heat}} \text{CH}_3\text{CONH}_2$ $\downarrow \text{Br}_2/\text{NaOH}$ CH_3NH_2 </div> <p>(ii)</p> <div style="border: 1px solid black; padding: 10px; margin: 10px auto; width: fit-content;"> $\text{CH}_3\text{-CH}_2\text{-C}\equiv\text{N} \xrightarrow{\text{H}_2/\text{Pt}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-NH}_2$ </div> <p>(or by any other method)</p> <p>b) Aniline undergoes resonance and as a result the electrons on the N-atom are less available for donation.</p> <p>c) (i) $(\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$ OR c) (ii) A = $\text{C}_6\text{H}_5\text{NH}_2$; B = $\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-$</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
30	<p>a) Due to presence of one unpaired electron in t_{2g} which gets excited to e_g / Due to excitation energy $t_{2g}^1 \rightarrow e_g^1$, it gives colour. (d-d transition) When heated, water is lost therefore crystal field splitting does not occur and it becomes colourless.</p> <p>b) The energy required to split the degenerate d-orbitals into two sets of orbitals (t_{2g} and e_g). / The difference of energy between the two sets of d-orbitals t_{2g} and e_g due to the presence of ligands in a definite geometry . OR b) (ii) $\Delta_o < P$, weak field ligand $\Delta_o > P$, strong field ligand</p> <p>c) Because the orbital splitting energies are not sufficiently large for forcing pairing / Due to low crystal field splitting energy.</p>	<p>1</p> <p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p>
	SECTION E	
31	<p>(a) (i)</p> $E_{\text{Cell}} = (E^{\circ}_c - E^{\circ}_a) - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{H}^+]^2}$ $= [(0) - (-0.76)] - \frac{0.059}{2} \log \frac{0.1}{(0.01)^2}$ $= 0.76 - 0.0295 \log 10^3$ $= 0.76 - 0.0885$ $= 0.6715 \text{ V} \quad \text{or } 0.67 \text{ V} \quad (\text{Deduct } \frac{1}{2} \text{ mark for no or incorrect unit})$ <p>(ii) The amounts of different substances liberated by the same quantity of electricity passing through the electrolytic solution are proportional to their chemical equivalent weights. 6F</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
	OR	
31	<p>(b) (i)</p> $\Lambda_m = \frac{\kappa}{c} \times 1000$ $\Lambda_m = \frac{2.48 \times 10^{-2}}{0.2} \times 1000$	<p>$\frac{1}{2}$</p>

	<p>(ii)</p> $\text{>C=O} \xrightarrow[\text{HCl}]{\text{Zn-Hg}} \text{>CH}_2 + \text{H}_2\text{O}$ <p>(I)</p> <p>(II)</p> $\text{C}_6\text{H}_5\text{CH}_3 + \text{CrO}_2\text{Cl}_2 \xrightarrow{\text{CS}_2} \text{C}_6\text{H}_5\text{CH(OCrOHCl}_2)_2 \xrightarrow{\text{H}_3\text{O}^+} \text{C}_6\text{H}_5\text{CHO}$	<p>1</p> <p>1</p>
	OR	
33	<p>(b)</p> <p>i)</p> <p>ii) Because semicarbazide undergoes resonance involving only one of the two - NH₂ groups, which is attached directly to the carbonyl-carbon atom.</p> <p>iii)</p> $\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{CrO}_3} 2 \text{ CH}_3\text{-CHO} \xrightleftharpoons{\text{dil. NaOH}} \text{CH}_3\text{-CH(OH)-CH}_2\text{-CHO}$ <p style="text-align: center;">Ethanal</p> <p>iv)</p> <p>v)</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>